



## 9. Poincaré group (20 points)

To be discussed on Wednesday, 17<sup>th</sup> December, 2025 in the tutorial.  
 Please indicate your preferences until Friday, 12/12/2025, 21:00:00 on the website.

### Exercise 9.1: Unitary representation of the Poincaré group

Let  $U(\Lambda, a)$  be a unitary representation of the Poincaré group with the multiplication law

$$(\Lambda_1, a_1)(\Lambda_2, a_2) = (\Lambda_1\Lambda_2, \Lambda_1a_2 + a_1),$$

where  $\Lambda \in \mathcal{L}_+^\uparrow$ ,  $\Lambda = (\Lambda^\mu{}_\nu)$  and  $a = (a^\mu)$ . Moreover, let

$$U_\rho(\omega, \epsilon) = e^{\frac{1}{2}\omega^{\mu\nu}M_{\mu\nu} + \epsilon^\mu P_\mu},$$

such that

$$\begin{aligned} U(\Lambda(\omega), 0) &= U_\rho(\omega, 0) \\ U(\mathbb{1}, a(\epsilon)) &= U_\rho(0, \epsilon), \end{aligned}$$

because

$$\Lambda \simeq \mathbb{1} + \omega + \mathcal{O}(\omega^2)$$

and also  $\omega^{\mu\nu} = -\omega^{\nu\mu}$ ,  $M_{\mu\nu} = -M_{\nu\mu}$ .

a) (3 points) Show that

$$U^{-1}(\Lambda, 0)U(\mathbb{1}, \epsilon)U(\Lambda, 0) = U(\mathbb{1}, \Lambda^{-1}\epsilon).$$

*Hint: Don't forget that the inverse  $U^{-1}(\Lambda, a)$  is also an element of the Poincaré group that shall satisfy*

$$U^{-1}(\Lambda, a)U(\Lambda, a) = U(\Lambda, a)U^{-1}(\Lambda, a) = U(\mathbb{1}, 0).$$

b) (3 points) Show that

$$U^{-1}(\Lambda, 0)P_\mu U(\Lambda, 0) = (\Lambda^{-1})^\nu{}_\mu P_\nu.$$

*Hint: Try to relate this with the previous task and  $U_\rho$ .*

c) (4 points) Find the commutator  $[M_{\mu\nu}, P_\rho]$ .

*Hint: Expand the relation of the previous task in 1st order in  $\omega$  and keep terms up to  $\omega$ . Also, keep in mind that a generic tensor  $T_{\mu\nu}$  can be written in terms of a symmetric and an antisymmetric component, that is*

$$T_{\mu\nu} = \frac{1}{2}(T_{\mu\nu} + T_{\nu\mu}) + \frac{1}{2}(T_{\mu\nu} - T_{\nu\mu}).$$

d) (3 points) Show that

$$U^{-1}(\Lambda, 0)U(\Lambda', 0)U(\Lambda, 0) = U(\Lambda^{-1}\Lambda'\Lambda, 0).$$

*Hint: Notice the similarity with task a).*

e) (4 points) Find the commutator  $[M_{\mu\nu}, M_{\rho\sigma}]$ .

*Hint: Expand the relation of the previous task in 1st order in  $\omega$ ,  $\omega'$  and keep terms up to  $\omega$ ,  $\omega'$  and  $\omega\omega'$ .*

f) (3 points) Show that  $[P_\mu, P_\nu] = 0$ .

*Hint: Try to do so via calculating  $[U(\mathbb{1}, a), U(\mathbb{1}, a')]$ .*