



5. Representation theory (16 points)

To be discussed on Tuesday, 2nd June, 2026 in the tutorial.

Please indicate your preferences until Thursday, 28/05/2026, 21:00:00 on the website.

Exercise 5.1: Killing form

During the lecture, we defined an inner product on the Lie algebra \mathfrak{g} called the *Killing form* κ . Let us study some of its properties.

- a) (2 points) An *automorphism* ω of \mathfrak{g} is a bijective map $\omega : \mathfrak{g} \rightarrow \mathfrak{g}$ which is compatible with the Lie bracket, i.e. it obeys

$$\omega([x, y]) = [\omega(x), \omega(y)].$$

Show that the Killing form κ is invariant under any automorphism of \mathfrak{g} .

- b) (2 points) Prove the ad-invariance of the Killing form, i.e. the property

$$\kappa([x, y], z) = \kappa(x, [y, z]), \quad \forall x, y, z \in \mathfrak{g}.$$

- c) (2 points) Let $\mathfrak{g} = \mathfrak{sl}(2, \mathbb{R}) = \text{span}(h, e, f)$ with Lie brackets $[h, e] = 2e$, $[h, f] = -2f$ and $[e, f] = h$. Compute the Killing form and show that it obeys

$$\kappa(x, y) = 4 \text{Tr}(xy), \quad \forall x, y \in \mathfrak{sl}(2, \mathbb{R}).$$

- d) (2 points) Repeat the exercise for $\mathfrak{so}(3)$. Is it semisimple?
e) (2 points) Repeat this for the three-dimensional Heisenberg algebra which we considered during the lecture. Is it semisimple?

Exercise 5.2: Low-dimensional Lie algebras

- a) (2 points) Consider the two-dimensional Lie algebra with generators T and U and Lie bracket $[T, U] = U$, and the three-dimensional Heisenberg algebra. Are these Lie algebras solvable? Are they nilpotent?
b) (2 points) Classify all two-dimensional Lie algebras up to isomorphism.

Exercise 5.3: Adjoint action

- a) (2 points) During the lecture, we defined the adjoint actions Ad and ad for Lie groups and Lie algebras respectively. Prove the identities

$$(\text{Ad}_g)^{-1} = \text{Ad}_{g^{-1}}, \quad \text{ad}_{\text{Ad}_g(X)} = \text{Ad}_g \text{ad}_X (\text{Ad}_g)^{-1}.$$