

Quantum Field Theory

lecturer: Falk Hassler (falk.hassler@uwr.edu.pl)

office: 448

lectures: Tue. 10:15 - 12:00 }
tutorials: Tue. 12:15 - 14:00 } 445

exercises & handwritten notes at

<https://www.fhassler.de/teaching>

- online ~ 1 week before tutorial
- assigned at Thur. 21:00 - u - \rightarrow email

Will be graded.

Problems? Contact me or course assistant


Biplab Mahato (334275@uwr.edu.pl)

exam

- in written @ end of semester
- at least 50% points of assigned exercise problems to qualify

office hours: Tue. 15:00 - 17:00

0. Motivation



1. Canonical quantisation

1.1. Klein-Gordon Field: Lagrangian

↗ It is governed by the action

$$S = \int d^4x \mathcal{L}(\phi, \partial_\mu \phi) \quad \text{Lagrangian}$$

$$\mathcal{L} = \frac{1}{2} \dot{\phi}^2 - \frac{1}{2} (\vec{\nabla} \phi)^2 - \frac{1}{2} m^2 \phi^2 \quad \dot{\phi} = \frac{\partial}{\partial t} \phi = \partial_t \phi$$

better in covariant form with metric $\vec{\nabla} = (\partial_{x^1}, \partial_{x^2}, \partial_{x^3})$

$$g_{\mu\nu} = g^{\mu\nu} = \begin{pmatrix} 1 & & & \\ & -1 & & \\ & & \dots & \\ 0 & & & -1 \end{pmatrix}$$

$$\boxed{\mathcal{L} = \frac{1}{2} (\partial_\mu \phi \partial^\mu \phi) - \frac{1}{2} m^2 \phi^2}$$

equation of motion from principle of least action

$$\delta S = 0$$

$$\delta S = \int d^4x \left[\frac{1}{2} \delta (\partial_\mu \phi \partial^\mu \phi) - \frac{1}{2} \delta (m^2 \phi^2) \right]$$

derivatives can be "removed" by integration by parts

$$\int d^4x \partial_\mu f_1 f_2 + \int d^4x f_1 \partial_\mu f_2 = \int d^4x \partial_\mu (f_1 f_2) = 0$$

we ignore boundary terms (at the moment)

$$\delta S = \int d^4x \left[-\partial_\mu \partial^\mu \phi - m^2 \phi \right] \delta \phi = 0$$

→

$$\boxed{\partial_\mu \partial^\mu \phi + m^2 \phi = 0 //}$$

Klein-Gordon equation

1.2. Hamiltonian Formulation

remember: classical mechanics positions q^i
conjugate momenta $p_i = \frac{\partial L}{\partial \dot{q}^i}$

Hamiltonian $H(t) = \sum_i p_i \dot{q}^i - L$ \Leftarrow Lagrangian

field theory $S = \int dt L = \int dt \int d^3x \mathcal{L}(\phi, \dot{\phi})$

$\pi(x) = \frac{\delta \mathcal{L}}{\delta \dot{\phi}(x)}$ conjugate momenta to ϕ

$H(t) = \int d^3x [\pi \dot{\phi} - \mathcal{L}] = \int d^3x \mathcal{H}$
Hamiltonian density

For the Klein-Gordon Lagrangian:

$$\pi(x) = \frac{\delta S}{\delta \dot{\phi}} \left(\frac{1}{2} \dot{\phi}^2 - \frac{1}{2} (\vec{\nabla} \phi)^2 - \frac{1}{2} m^2 \phi^2 \right) = \dot{\phi}$$

$$H = \int d^3x \left[\frac{1}{2} \pi^2 + \frac{1}{2} |\nabla \phi|^2 + \frac{1}{2} m^2 \phi^2 \right]$$

Field equations:

- | |
|-----------------------------|
| 1.) Hamiltonian ✓ |
| 2.) Poisson brackets (Pb's) |
| 3.) Time evolution |

2.) $\{ \underbrace{\phi(t, \vec{x})}_{\text{equal time}}, \underbrace{\pi(t, \vec{y})}_{\text{equal time}} \} = \delta(\vec{x} - \vec{y})$

all other Pb's are 0

3.) for all functions of ϕ and π , $O(\phi, \pi)$, we have

$$\frac{\partial}{\partial t} \Theta = \{ \Theta, H \}$$

Let's check for
 $\Theta(\phi, \pi) = \phi$:

$$\begin{aligned} \frac{\partial}{\partial t} \phi(t, \vec{x}) &= \int d^3y \{ \phi(t, \vec{x}), 1/2 \pi^2(t, \vec{y}) \} \\ &= \int d^3y \underbrace{\{ \phi(t, \vec{x}), \pi(t, \vec{y}) \}}_{\delta(\vec{x} - \vec{y})} \pi(t, \vec{y}) \\ &= \pi(t, \vec{x}) \end{aligned}$$

$$\frac{\partial}{\partial t} \pi(t, \vec{x}) = \dots = (\vec{\nabla}^2 - m^2) \phi(t, \vec{x})$$

$$\dot{\phi} = \pi = (\vec{\nabla}^2 - m^2) \phi$$

$$\ddot{\phi} - \vec{\nabla}^2 \phi + m^2 \phi = \underline{\underline{\partial_\mu \partial^\mu \phi + m^2 \phi = 0}}$$

Klein-Gordon equation

1.3. Quantisation

Fourier transformation \rightarrow momentum space

$$\phi(t, \vec{x}) = \int \frac{d^3p}{(2\pi)^3} e^{i\vec{x} \cdot \vec{p}} \phi(t, \vec{p})$$

$$\text{Klein-Gordon eq. } \left[\frac{\partial^2}{\partial t^2} + \underbrace{(|\vec{p}|^2 + m^2)}_{\omega_{\vec{p}}^2} \right] \phi(t, \vec{p}) = 0$$

$$\omega_{\vec{p}} = \sqrt{|\vec{p}|^2 + m^2}$$

\hookrightarrow Harmonic oscillator with frequency $\omega_{\vec{p}}$

first one HO

$$\left. \begin{aligned} H_{HO} &= \frac{1}{2} p^2 + \frac{1}{2} \omega^2 \phi^2, & \phi &= \frac{1}{\sqrt{2\omega}} (a + a^\dagger) \\ p &= -i \sqrt{\frac{\omega}{2}} (a - a^\dagger) \end{aligned} \right\} \text{with } [\phi, p] = i\hbar$$

implies $[a, a^\dagger] = 1$
↖ raising operator
↙ lowering operator

Hilbert space: $a|0\rangle = 0$ vacuum or ground state
 $(a^\dagger)^n |0\rangle = |n\rangle$

→ $H_{H_0} = \omega (a^\dagger a + 1/2)$ $a^\dagger a |n\rangle = n |n\rangle$
 $H_{H_0} |n\rangle = \omega(n + 1/2) |n\rangle$

next KG theory

usually suppressed

$$\phi(t, \vec{x}) = \int \frac{d^3 p}{(2\pi)^3} \frac{1}{\sqrt{2\omega_{\vec{p}}}} (a_{\vec{p}}(t) + a_{-\vec{p}}^\dagger(t)) e^{i\vec{p}\cdot\vec{x}}$$

$$\pi(t, \vec{x}) = \int \frac{d^3 p}{(2\pi)^3} (-i) \sqrt{\frac{\omega_{\vec{p}}}{2}} (a_{\vec{p}} - a_{-\vec{p}}^\dagger) e^{i\vec{p}\cdot\vec{x}}$$

with $[a_{\vec{p}}, a_{\vec{p}'}^\dagger] = (2\pi)^3 \delta(\vec{p} - \vec{p}')$ from
 $[\phi(\vec{x}), \pi(\vec{y})] = i \delta(\vec{x} - \vec{y})$ (please check)

Comparing with $\{\phi(\vec{x}), \pi(\vec{y})\} = \delta(\vec{x} - \vec{y})$ we find

$\{., .\} \rightarrow i\hbar [., .]$ <small>"1 for us"</small>	<u>Canonical Quantisation</u>
---	-----------------------------------

$$H = \int \frac{d^3 p}{(2\pi)^3} \omega_{\vec{p}} (a_{\vec{p}}^\dagger a_{\vec{p}} + \underbrace{1/2 [a_{\vec{p}}, a_{\vec{p}}^\dagger]}_{\text{vacuum energy}})$$

Spectrum: ↖ vacuum state ∞ = vacuum energy

$$a_{\vec{p}} |0\rangle = 0$$

$$a_{\vec{p}}^\dagger |0\rangle = 1\text{-particle state with momentum } \vec{p}$$

and energy $E_{\vec{p}} = \omega_{\vec{p}} = \sqrt{|\vec{p}|^2 + m^2}$
 (remember $c=1$)

1.4. Heisenberg picture & propagator

$$\phi(x) = e^{iHt} \phi(\vec{x}) e^{-iHt}$$

4-position $x^\mu = (x^0, \underbrace{x^1, x^2, x^3}_{\vec{x}})$

same for $\pi(x)$

now we have: $i \frac{\partial}{\partial t} \phi = [\phi, H]$

(compare with) $\frac{\partial}{\partial t} \phi = \{ \phi, H \}$

$$\phi(x) = \int \frac{d^3p}{(2\pi)^3} \frac{1}{\sqrt{2E_{\vec{p}}}} \left(a_{\vec{p}} e^{-i p_\mu x^\mu} + a_{\vec{p}}^\dagger e^{i p_\mu x^\mu} \right) \Big|_{p^0 = E_{\vec{p}}}$$

on-shell \rightarrow

$$\pi(x) = \frac{\partial}{\partial t} \phi(x) = \dot{\phi}$$


Propagator:

EXPERIMENT

create particle at position y and
 detect it at position x

$$D(x-y) = \langle 0 | \phi(x) \phi(y) | 0 \rangle$$

$$= \int \frac{d^3p}{(2\pi)^3} \frac{1}{2E_{\vec{p}}} e^{-ip(x-y)}$$




$$D_F(x-y) = \Theta(x^0 - y^0) \langle 0 | \phi(x) \phi(y) | 0 \rangle +$$

\mathbb{R} Feynman

$$\Theta(y^0 - x^0) \langle 0 | \phi(y) \phi(x) | 0 \rangle$$

$$\Theta(x) = \begin{cases} 1 & x > 0 \\ 0 & x \leq 0 \end{cases}$$



$$= \lim_{\epsilon \rightarrow 0^+} \mp \frac{1}{2\pi i} \int_{-\infty}^{\infty} \frac{1}{\tau \pm i\epsilon} e^{\mp i x \tau} d\tau \quad \rightarrow \text{EX 2.1.}$$

$$D_F(x-y) = \lim_{\epsilon \rightarrow 0^+} \int \frac{d^4 p}{(2\pi)^4} \frac{i}{p^2 - m^2 + i\epsilon} e^{-i p(x-y)}$$

Feynman propagator for KG field

$$D_F(x-y) = \Theta(x^0 - y^0) D(x-y) + \Theta(y^0 - x^0) D(y-x)$$

$$= \langle 0 | T \phi(x) \phi(y) | 0 \rangle$$

↖ Time ordering operators

"later" operators go to the left

2. Dirac Field

2.1. Lorentz transformations

elements of Lorentz group $O(3,1)$

coordinates: $x^\mu \rightarrow x'^\mu = \Lambda^\mu_\nu x^\nu$

such that $x^\mu x^\nu g_{\mu\nu} = x'^\mu x'^\nu g_{\mu\nu}$

Scalar fields: $\phi(x) \rightarrow \phi'(x) = \phi(\Lambda^{-1}x)$

Vector fields: $V^\mu(x) \rightarrow V'^\mu(x) = \Lambda^\mu_\nu V^\nu(\Lambda^{-1}x)$

example $\partial^\mu \phi(x) = V^\mu(x)$

dual, 1-form: $A_\mu(x) \rightarrow A'_\mu(x) = (\Lambda^{-1})^\nu_\mu A_\nu(\Lambda^{-1}x)$

fields

then $A_\mu V^\mu$ is a scalar

example $A_\mu(x) = \partial_\mu \phi(x)$

- Questions:
- Are there more examples? Yes there are!
 - How do we classify them?

↗ Monographic lecture "Lie algebras & Lie groups"

$so(3,1)$ Lie algebra

$\frac{1}{2} \cdot 4(4-1) = 6$ generators $J^{\mu\nu} = -J^{\nu\mu}$

defined by commutators:

$$[J^{\mu\nu}, J^{\alpha\beta}] = i(g^{\nu\beta} J^{\mu\alpha} - g^{\mu\beta} J^{\nu\alpha} - g^{\nu\alpha} J^{\mu\beta} + g^{\mu\alpha} J^{\nu\beta})$$

1) $J_1 = J^{23}$, $J_2 = J^{31}$, $J_3 = J^{12}$ generate rotations of 3 spacial dir.

2) $K_1 = J^{01}$, $K_2 = J^{02}$, $K_3 = J^{03}$ boosts

$$[J_i, J_j] = i \sum_{k=1}^3 \epsilon_{ijk} J_k = i \epsilon_{ijk} J_k \quad so(3) \text{ sub algebra}$$

$$[J_i, K_j] = i \epsilon_{ijk} K_k \quad \text{and} \quad [K_i, K_j] = -i \epsilon_{ijk} J_k$$

Task: Find explicit representations for matrices $J^{\mu\nu}$!

2.2. γ -matrices and the Dirac algebra

More concret: find 4 $n \times n$ matrices γ^μ with

$$\{\gamma^\mu, \gamma^\nu\} = \gamma^\mu \gamma^\nu + \gamma^\nu \gamma^\mu = 2 g^{\mu\nu} \cdot \mathbb{1}_{n \times n} \leftarrow n \times n \text{ identity matrix}$$

Ex 1.2 check that they generate $so(3,1)$!

$$\gamma^0 = \begin{pmatrix} 0 & \mathbb{1}_{2 \times 2} \\ \mathbb{1}_{2 \times 2} & 0 \end{pmatrix}, \quad \gamma^i = \begin{pmatrix} 0 & \sigma^i \\ \sigma^i & 0 \end{pmatrix} \quad \sigma^i = \text{Pauli matrices}$$

$$\sigma^1 = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \quad \sigma^2 = \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix}, \quad \sigma^3 = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$$

We also need one more γ -matrix:

$$\gamma^5 = i \gamma^0 \gamma^1 \gamma^2 \gamma^3 = \begin{pmatrix} -\mathbb{1}_{2 \times 2} & 0 \\ 0 & \mathbb{1}_{2 \times 2} \end{pmatrix} \quad \text{with} \quad \{\gamma^5, \gamma^\mu\} = 0$$

and therefore $[\gamma^5, J^{\mu\nu}] = 0$

γ -matrices act on 4-component vectors

$$\Psi = \begin{pmatrix} \Psi_L \\ \Psi_R \end{pmatrix} \quad \text{called} \quad \underline{\underline{\text{Dirac-spinors}}}$$

They decompose into 2 fundamental irreps of $so(3,1)$, the 2-component (but complex) Weyl spinors Ψ_L & Ψ_R .

They are the ± 1 eigenvalues of γ^5 .

Contraction of two Dirac-spinors \rightarrow Lorentz scalar

naively: $\Psi^\dagger \Psi$ (like for vector) does not work!

rather $\bar{\Psi} \Psi$ with

$$\boxed{\bar{\Psi} = \Psi^\dagger \gamma^0} \quad \text{Dirac conjugation}$$

EX 1.2 verify that $\bar{\Psi} \Psi$ is a Lorentz scalar

2.3. Dirac equation

$$\boxed{S_{\text{Dirac}} = \int d^4x \bar{\Psi} (i \gamma^\mu \partial_\mu - m) \Psi}$$

notation $\gamma^\mu \partial_\mu = \not{\partial}$
or $\gamma^\mu p_\mu = \not{p}$

field equation:
$$\frac{\delta S_{\text{Dirac}}}{\delta \bar{\Psi}} = 0$$

results in
$$\boxed{(i \not{\partial} - m) \Psi = 0} = \text{Dirac equation}$$

plane wave solutions

$$\Psi(x) = u(p) e^{-i p x} + v(p) e^{i p x}, \quad p^0 > 0$$

↳ $(\not{p} - m) u(p) = 0$ and $(\not{p} + m) v(p) = 0$

both have two linearly independent solutions

$$u(p) = u^s(p) \quad s=1,2 \quad \text{and} \quad v(p) = v^r(p) \quad r=1,2$$

which can be normalised to

$$\begin{aligned} \bar{u}^r(p) u^s(p) &= 2m \delta^{rs} & \bar{u}^r(p) v^s(p) &= 0 \\ \bar{v}^r(p) v^s(p) &= -2m \delta^{rs} & \bar{v}^r(p) u^s(p) &= 0 \end{aligned}$$

Interpretation of them as electrons and positrons
(see EX 1.3)

2.4. Quantisation of the Dirac field

conjugate momentum to Ψ is $i \Psi^\dagger$

Hamiltonian:
$$H = \int d^3x \bar{\Psi} (-i \vec{\gamma} \cdot \vec{\nabla} + m) \Psi$$

spacial part $\vec{\nabla}$ only

mode expansion:

$$\Psi(x) = \int \frac{d^3p}{(2\pi)^3} \frac{1}{\sqrt{2E_{\vec{p}}}} \sum_s \left(a_{\vec{p}}^s u^s(p) e^{-ipx} + (b_{\vec{p}}^s)^\dagger v^s(p) e^{ipx} \right)$$

$$\bar{\Psi}(x) = \int \frac{d^3p}{(2\pi)^3} \frac{1}{\sqrt{2E_{\vec{p}}}} \sum_s \left(b_{\vec{p}}^s \bar{v}^s(p) e^{-ipx} + (a_{\vec{p}}^s)^\dagger \bar{u}^s(p) e^{ipx} \right)$$

$$\{a_{\vec{p}}^r, (a_{\vec{q}}^s)^\dagger\} = \{b_{\vec{p}}^r, (b_{\vec{q}}^s)^\dagger\} = (2\pi)^3 \delta(\vec{p}-\vec{q}) \delta^{rs}$$

↑ ↙ not Poisson brackets, but anti-commutator!
↘ $\{a, b\} = a \cdot b + b \cdot a$

all other $\{.,.\} = 0$

reason for $\{.,.\}$ instead of $[.,.]$ is that we are dealing with fermions

Vacuum $|0\rangle$ annihilated by

$$a_{\vec{p}}^s |0\rangle = b_{\vec{p}}^s |0\rangle = 0$$

We can only have one particle with given state

$$(a_{\vec{p}}^1)^\dagger |0\rangle \quad \text{because} \quad (a_{\vec{p}}^1)^\dagger (a_{\vec{p}}^1)^\dagger |0\rangle = 0$$
$$\{a_{\vec{p}}^1, a_{\vec{p}}^1\} = 2(a_{\vec{p}}^1)^2 = 0$$

= Pauli exclusion principle!

$$\text{Hamiltonian: } H = \int \frac{d^3p}{(2\pi)^3} \sum_s E_{\vec{p}} \left(a_{\vec{p}}^{s\dagger} a_{\vec{p}}^s + b_{\vec{p}}^{s\dagger} b_{\vec{p}}^s \right)$$

and Feynman propagator:

$$D_F(x-y) = \langle 0 | T \Psi(x) \bar{\Psi}(y) | 0 \rangle$$
$$= \int \frac{d^4p}{(2\pi)^4} \frac{i(\not{p} + m)}{p^2 - m^2 + i\epsilon} e^{-ip(x-y)}$$