

## 5.6. Quantisation of Fermions

### 5.6.1. Grassmann numbers

Remember 2.4.: for Dirac field  $\{\hat{a}_{\vec{p}}^r, \hat{a}_{\vec{q}}^s\} = (2\pi)^3 \delta(\vec{p}-\vec{q}) \delta^{rs}$   
 anti-commutator

Fine for operators, but in path integral we have just ordinary numbers (like  $\hat{\phi}$  vs.  $\phi$  from last two lectures).

Question: What shall we do there to implement fermionic fields?

Answer: Grassmann numbers:

$$\Theta\eta = -\eta\Theta$$

implies that  $\Theta^2 = 0$  and that any function  $f(\Theta)$  can be written as

$$f(\Theta) = A + \Theta B \quad (\text{Taylor expansion})$$

We further need: 1) Integration  $\int d\Theta f(\Theta)$

$= \int d\Theta (A + \overset{\circ}{B}\Theta)$  should be invariant under

$$= \int d\Theta ([A + B\eta] + \overset{\circ}{B}\Theta) \stackrel{\Theta \rightarrow \Theta + \eta}{=} B$$

$$\boxed{\int d\Theta (A + B\Theta) = B}$$

multiple integrals:  $\int d\Theta \int d\eta \eta \Theta = 1$

$$2) \text{ Differentiation } \frac{\partial}{\partial \Theta} f(\Theta) = \frac{\partial}{\partial \Theta} (A + B\Theta) = B$$

for Grassmann numbers integration = differentiation

3) (complex Grassmann numbers)

$$\Theta = \frac{\Theta_1 + i\Theta_2}{\sqrt{2}}, \quad \Theta^* = \frac{\Theta_1 - i\Theta_2}{\sqrt{2}}$$

$$4) \text{ complex conjugation } (\theta \eta)^* = \eta^* \theta^* = -\theta^* \eta^*$$

we can now evaluate the Grassmann version of a Gaussian integral

$$\begin{aligned} I &= \int d\theta^* d\theta e^{-\theta^* b \theta} \xrightarrow{\text{Taylor expand}} \int d\theta^* d\theta (1 - \theta^* b \theta) \\ &= \int d\theta^* d\theta (1 + \theta b \theta^*) = b // \end{aligned}$$

remember: if  $\theta$  would be  $\in \mathbb{C}$ ,  $I = \frac{2\pi}{b}$

$$\int d\theta^* d\theta \theta \theta^* e^{-\theta^* b \theta} = 1 = \frac{1}{b} \cdot b$$

now in  $N$  dimensions  $I = \left( \prod_i \int d\theta_i^* d\theta_i \right) e^{-\sum_i \theta_i^* B_{ij} \theta_j}$

I) diagonalise  $B_{ij}$  with unitary transformation

$$\theta_i' = U_{ij} \theta_j$$

$$\prod_i \theta_i' = \frac{1}{N!} \epsilon^{i_1 i_2 \dots i_N} \theta_1' \theta_2' \dots \theta_N' = \dots = (\det U) \left( \prod_i \theta_i \right)$$

$$\Rightarrow I = \left( \prod_i \int d\theta_i'^* d\theta_i' \right) e^{-\sum_i \theta_i'^* b_i \theta_i} = \prod_i b_i = \det B //$$

$$\boxed{\left( \prod_i \int d\theta_i^* d\theta_i \right) e^{-\sum_i \theta_i^* B_{ij} \theta_j} = \det B}$$

### 5.6.2. Dirac Propagator

remember Dirac spinor  $\hat{\Psi} = (\hat{\psi}_1 \ \hat{\psi}_2 \ \hat{\psi}_3 \ \hat{\psi}_4)^T$ ; now it becomes  $\Psi(x) = \sum_i \Psi_i \phi_i(x)$

Grassmann number  $\nearrow$  ordinary Dirac spinor basis

$$\text{i.e.: } \phi_1 = \begin{pmatrix} 1 \\ 0 \\ 0 \\ 0 \end{pmatrix}, \dots, \phi_4 = \begin{pmatrix} 0 \\ 0 \\ 0 \\ 1 \end{pmatrix}$$

we can now evaluate:

$$\langle 0 | T \psi(x_1) \bar{\psi}(x_2) | 0 \rangle = \frac{\int D\bar{\psi} D\psi e^{iS} \psi(x_1) \bar{\psi}(x_2)}{\int D\bar{\psi} D\psi e^{iS}}$$

with  $S = \int d^4x \bar{\psi}(i\gamma^\mu) \psi$

$$\langle 0 | T \psi(x_1) \bar{\psi}(x_2) | 0 \rangle = S_F(x_1 - x_2) = \int \frac{d^4k}{(2\pi)^4} \frac{i e^{-ik(x_1 - x_2)}}{k - m + i\epsilon}$$

Note: higher correlation functions are obtained from Wick's theorem

### 5.6.3. Generating Function

$$Z[\bar{\eta}, \eta] = \int \mathcal{D}\bar{\psi} \mathcal{D}\psi \exp [iS + \int d^4x [\bar{\eta}\psi + \bar{\psi}\eta]]$$

Grassmann valued source field

$$= Z_0 \exp \left[ - \int d^4x \int d^4y \bar{\eta}(x) S_F(x-y) \eta(y) \right]$$

and therefore

$$\langle 0 | T \psi(x_1) \bar{\psi}(x_2) | 0 \rangle = Z_0^{-1} \left( -i \frac{\delta}{\delta \bar{\eta}(x_1)} \right) \left( i \frac{\delta}{\delta \eta(x_2)} \right) Z[\bar{\eta}, \eta] \Big|_{\bar{\eta}, \eta = 0}$$

### 5.7. Symmetries in the Path Integral

three point correlator for free scalar theory

$$\langle 0 | T \phi_1 \phi_2 \phi_3 | 0 \rangle = Z_0^{-1} \int \mathcal{D}\phi e^{iS} \phi_1 \phi_2 \phi_3$$

$$S = i \int dx \left[ \frac{1}{2} (\partial_\mu \phi)^2 - \frac{1}{2} m^2 \phi^2 \right]$$

now shift  $\phi(x) \rightarrow \phi'(x) = \phi(x) + \varepsilon(x)$ ; leaves  $\mathcal{D}\phi$  invariant,  $\mathcal{D}\phi = \mathcal{D}\phi'$

$$0 = \int \mathcal{D}\phi e^{iS} \left[ i \int d^4x \varepsilon (-\partial^2 - m^2) \phi \cdot \phi_1 \phi_2 \phi_3 + \varepsilon_1 \phi_1 \phi_2 \phi_3 + \phi_1 \varepsilon_2 \phi_3 + \phi_1 \phi_2 \varepsilon_3 \right]$$

$$(\partial^2 + m^2) \langle 0 | T \phi \phi_1 \phi_2 \phi_3 | 0 \rangle = -i \delta(x-x_1) \langle 0 | T \phi_2 \phi_3 | 0 \rangle + \text{cycl.}$$

in particular:  $(\partial^2 + m^2) \langle 0 | T \phi \phi_1 | 0 \rangle = -i \delta(x-x_1)$

We see: Feynman propagator is the Green's function of the classical field equations.

This idea generalises to arbitrary Lagrangians  $\mathcal{L}$  (as long as the measure is preserved under shifts)

$$\left\langle \left[ \frac{\partial \mathcal{L}}{\partial \phi} - \partial_\mu \left( \frac{\partial \mathcal{L}}{\partial (\partial_\mu \phi)} \right) \right] \phi_1 \dots \phi_n \right\rangle = \sum_{i=1}^n \left\langle \phi_1 \dots (i \delta(x-x_i)) \dots \phi_n \right\rangle$$

Schwinger - Dyson equation

contact term